

NOTES ON THE ECOLOGY OF FOUR SPECIES OF  
PYGOPODID LIZARDS IN THE  
GREAT VICTORIA DESERT

By ERIC R. PIANKA  
Section of Integrative Biology C0930,  
University of Texas at Austin,  
Austin, Texas 78712 USA

ABSTRACT

Following a brief review of the literature on pygopodid lizards, ecological data are presented for *Delma butleri*, *Delma nasuta*, *Lialis burtonis*, and *Pygopus nigriceps*. *D. nasuta* and *D. butleri* are both diurnal and nocturnal, *L. burtonis* is crepuscular, and *P. nigriceps* is nocturnal. Active body temperatures were highest in *D. butleri* and lowest in *P. nigriceps*. Both *Delma* species climb inside spinifex tussocks, *L. burtonis* and *P. nigriceps* are terrestrial, but one *P. nigriceps* was found 0.2m above ground on top of a spinifex tussock. Most individuals of both species of *Delma* were found on desert flats. *Lialis burtonis* and *P. nigriceps* had broader habitat niche breadths and were associated with both flats and sand ridges. Tails are shortest in *L. burtonis* and *P. nigriceps*, and 3+ times snout-vent length in both *Delma* species. All these pygopodids are dietary specialists, although food niche breadths vary from 1.0 to 4.26. *Lialis burtonis* is most specialized, eating only lizards and largely skinks. Mating occurs during the Austral Spring, clutch sizes always consist of 2 eggs and relative clutch mass varied from 0.137 to 0.176 among species.

INTRODUCTION

Pygopodids are uncommon Australian lizards, sometimes called flap-foots due to their complete lack of fore limbs and greatly reduced vestigial hind limbs. Pygopodids, which have undergone an adaptive radiation, are a monophyletic family derived from Diplodactylid

geckos (Kluge 1974, 1976; Jennings *et al.* 2003). These cryptic lizards are difficult to find and study and relatively little is known about most of the 40-odd described species. Six genera are recognized, two of which (*Ophidiocephalus* and *Pletholax*), are monotypic and scarcely known (but see Downes *et al.* 1997; Ehmann 1978, 1981, 1992;

and Bamford 1998). *Delma* (including the former "Aclys") is the most species-rich genus, with more than 20 named widespread species, which include both terrestrial and arboreal species. Some *Delma* can lift themselves completely off the ground using their long and powerful tails (Bauer 1986). Most "swim" rapidly through grass tussocks. The second largest genus *Aprasia* is also widespread, with 11 species, all of which are burrowers with relatively short tails (Parker 1956). *Aprasia* have converged on blind snakes; their diets consist of larvae and pupae of ants of several genera (Webb and Shine 1994). Food habits and reproductive biology of pygopodids were reviewed by Patchell and Shine (1986). *Pygopus* (including "Paradelma") includes 5 terrestrial nocturnal species that mimic venomous snakes. These lizards rear up their heads and feint menacingly, hissing and lunging like snakes, but give away their true identity when they flick out their short black blunt gekkonid tongue. One of the best studied species is *Lialis burtonis*, an ecological equivalent of a snake that sits and waits for large skinks and has hinged teeth like some skink-eating snakes (Savitzky 1981; Patchell and Shine 1986a, 1986b, 2009). As a further example of its convergence on snakes, caudal luring has been reported in *Lialis* (Murray *et al.* 1991). Moreover, *Lialis* has elliptical pupils and is crepuscular as are many snakes (Neill 1957). Evolutionary advantages of limb-

lessness in pygopodids are discussed by Shine (1986).

## METHODS

Over the last 42 years, from September 1966 through November 2008, on 11 separate research expeditions, I have spent 41 months and 1256 days in the field studying lizards, mostly during Austral Springs in the Great Victoria Desert (GVD) of Western Australia. However, a few specimens were captured outside the GVD. Up until 1979, all lizards were collected by hand, and data were obtained on date and time of activity, ambient air temperature, active body temperature, habitat and micro-habitat. Data for 3 species of pygopodids were summarized in Pianka (1986), in which *D. butleri* was unfortunately misidentified as "*D. fraseri*." Beginning in 1989, most lizards were collected using pit traps (62,226 pit trap days), which provided more limited qualitatively different information.

Items within stomachs were sorted among 15 categories, mostly arthropod orders. Items were counted and volumes estimated to the nearest cubic millimetre for each category. Volumes were estimated by placing a one millimetre thick layer of material over square millimetre grid paper and approximating total volume. Each lizard's counted stomach contents were kept individually and stored in ethanol. Dietary niche breadths were estimated

using the inverse of Simpson's index of diversity [ $D = 1 / \sum p_i^2$ ] where  $p_i$  is the proportion by volume of food items in stomachs based on 15 prey categories.

All lizards were collected and deposited in the LA County Museum of Natural History and the Western Australian Museum. Here, I report augmented ecological information on 4 species of pygopodids: *Delma butleri* (N=101), *Delma nasuta* (N=15), *Lialis burtonis* (N=60), and *Pygopus nigriceps* (N=57).

## RESULTS

### Time of Activity

Most pygopodids were active during the warmer months of October through February. In the following, all times are decimalized in metric units. *Delma butleri* are active during the day (mean time of day = 10.02, N=12, range 7.5 to 12.42), but also in late afternoon and evening hours (mean time = 18.76, N=6, range 16.62 to 20.58). Of the 15 *Delma nasuta* collected, one was active at 8.5 (0830) AM, 11 were captured or pit trapped during the night or in early AM, two were active

during daylight hours and pit trapped during the afternoon, and one was found active at night. Three *Lialis* were active during morning hours (mean time 8.5, range 7.3 to 9.12) and 9 were active in the afternoon and evening (mean = 16.87, range 14.6 to 19.5), indicating crepuscular activity. All *Pygopus* were active at night (mean time = 19.72, N=16, range 19 to 21.57).

### Temperature Relationships

In this study (Table 1), the diurnal *Delma butleri* had the highest active body temperatures ranging from 27.7 to 35.2°C, with a mean of 31.74°C. Air and body temperature was obtained for only one *D. nasuta* which was active at 8.5 (0830) AM (air = 24.4 and body = 29.8°C). Average body temperature chosen by *Lialis* in a laboratory thermal gradient was 35.1°C (Bradshaw *et al.* 1980); higher body temperatures were selected after feeding and lower body temperatures when lizards were starved. Body temperatures of eleven wild-caught individual *Lialis* ranged from 23.2 to 37.7°C with a mean of 29.0°C. Active body temperatures of nocturnal *Pygopus* were lower, ranging from

Table 1. Ambient air temperatures and active body temperatures for 3 species with data, along with the slope of regressions of active body temperatures on ambient air temperatures (Sample sizes in parentheses).

| Species             | Air Temperature (N) | Body Temperature (N) | Slope |
|---------------------|---------------------|----------------------|-------|
| <i>D. butleri</i>   | 28.48 (16)          | 31.74 (8)            | 0.560 |
| <i>L. burtonis</i>  | 25.9 (12)           | 29.0 (11)            | 0.789 |
| <i>P. nigriceps</i> | 24.42 (15)          | 24.88 (12)           | 0.762 |

20.8 to 28.2 (mean 24.88). Slopes of regressions of active body temperature on ambient air temperature were lower for *Delma butleri* an indication of thermoregulation (Huey and Slatkin 1976). Steeper slopes for *Lialis* and *Pygopus*, relative to *Delma*, suggest that they are thermoconformers.

#### Microhabitat and Habitat

Both species of *Delma* were strongly associated with spinifex grass (*Triodia basedowi*), through which they "swim" rapidly and with ease. All *Lialis* were terrestrial and were found in a variety of microhabitats. *Pygopus* are largely terrestrial although one individual was active 0.2 metres above ground on top of a spinifex tussock. Most *D. butleri*

and *D. nasuta* were found on desert flats. *Lialis* and *Pygopus* were distributed more evenly across habitats, being associated with the flats as well as sandridges, and have broader habitat niche breadths (Table 2).

#### Size, Relative Tail Length

*Pygopus* and *Lialis* are large and have relatively short tails, whereas both species of *Delma* are smaller and have considerably longer tails, relative to their snout-vent lengths (Table 3).

#### Diets

All these pygopodids are dietary specialists, although food niche breadths calculated with Simpson's index of diversity (see above) vary from 1.0 to 4.26 (Table 4). *Lialis* is most specialized,

Table 2. Percentages of lizards found in various habitats (Sample sizes in parentheses). Lizards found at an interface between habitats are split between both. Habitat niche breadths calculated with the inverse of Simpson's index of diversity,  $1/\sum p_i^2$  where  $p_i$  represents the proportion of animals in habitat  $i$ .

| Species             | Flat      | Base        | Slope      | Crest       | Total | Habitat NB |
|---------------------|-----------|-------------|------------|-------------|-------|------------|
| <i>D. butleri</i>   | 84.2 (85) | 14.9 (15)   | 0.5 (0.5)  | 0.5 (0.5)   | 101   | 1.37       |
| <i>D. nasuta</i>    | 66.7 (10) | 13.3 (2)    | 6.7 (1)    | 13.3 (2)    | 15    | 2.06       |
| <i>L. burtonis</i>  | 35.0 (21) | 29.2 (17.5) | 15.8 (9.5) | 20.0 (12)   | 60    | 3.67       |
| <i>P. nigriceps</i> | 31.6 (18) | 36.8 (21)   | 7.9 (4.5)  | 23.7 (13.5) | 57    | 3.36       |

Table 3. Average snout-vent length (SVL) and tail lengths (in mm), ratio of tail length over SVL, and mean body weight (in grams).

| Species             | Snout-Vent Length (N) | Tail Length (N) | Tail/SVL (N) | Weight (N) |
|---------------------|-----------------------|-----------------|--------------|------------|
| <i>D. butleri</i>   | 67.9 (99)             | 200.1 (58)      | 3.10 (58)    | 3.34 (98)  |
| <i>D. nasuta</i>    | 86.9 (15)             | 261.3 (7)       | 3.19 (7)     | 6.45 (15)  |
| <i>L. burtonis</i>  | 156 (44)              | 186 (23)        | 1.27 (23)    | 9.89 (42)  |
| <i>P. nigriceps</i> | 160 (58)              | 217 (27)        | 1.42 (27)    | 13.46 (58) |

Table 4. Proportions of 15 different prey categories by volume in pygopodid stomachs.

| Species/Prey Type            | <i>D. butleri</i> | <i>D. nasuta</i> | <i>Lialis</i> | <i>Pygopus</i> |
|------------------------------|-------------------|------------------|---------------|----------------|
| Centipedes                   | 0.0435            | 0                | 0             | 0              |
| Spiders                      | 0.3897            | 0.0787           | 0             | 0.2681         |
| Scorpions                    | 0                 | 0                | 0             | 0.2691         |
| Ants                         | 0.0035            | 0                | 0             | 0.0165         |
| Orthopterans                 | 0.2609            | 0.7191           | 0             | 0.1094         |
| Thysanura                    | 0.0486            | 0.0224           | 0             | 0              |
| Blattaria                    | 0.0400            | 0                | 0             | 0              |
| Phasmids/Mantids             | 0.0052            | 0                | 0             | 0              |
| Isoptera                     | 0.0626            | 0.0337           | 0             | 0.0202         |
| Diptera                      | 0.0209            | 0                | 0             | 0.0101         |
| Larvae                       | 0.0453            | 0                | 0             | 0              |
| Other Insects                | 0.0261            | 0.1011           | 0             | 0              |
| Vertebrates                  | 0                 | 0                | 1.0           | 0              |
| Unidentified                 | 0.0208            | 0                | 0             | 0.2965         |
| Other                        | 0                 | 0.0449           | 0             | 0.0101         |
| Total Volume of Prey, cc.    | 5.75              | 0.89             | 10.75         | 5.94           |
| Dietary Niche Breadth        | 4.26              | 1.86             | 1.0           | 3.94           |
| Number of Stomachs           | 85                | 14               | 44            | 49             |
| Percentage of Empty Stomachs | 29.1              | 35.7             | 79.5          | 36.7           |

eating only other lizards, largely skinks (80% of stomachs were empty). Species of skinks found in *Lialis* stomachs include *Lerista bipes*, *Menetia greyi*, *Ctenotus dux*, and *Ctenotus quattuordecimlineatus*. *Pygopus* eat predominantly spiders and scorpions, which they capture when these arthropods are active above ground at night. *Delma butleri* consume spiders and orthopterans, and *Delma nasuta* prey mainly on orthopterans.

### Reproduction

In all 4 species, testes were enlarged in September, October and November suggesting that mating takes place during the Austral Spring. Two *Delma butleri* females each contained 2 eggs in

their oviducts, clutch volume as a fraction of female body weight (relative clutch mass, RCM) were 0.132 and 0.160 (mean 0.146). Four *Lialis* females each had 2 eggs in their oviducts, clutch volumes as a fraction of body weight for these females were 0.114, 0.130, 0.150 and 0.153 (mean RCM 0.137). Two *Pygopus nigriceps* females each had 2 eggs in their oviducts, with relative clutch masses of 0.116 and 0.235 (mean RCM 0.176).

### ACKNOWLEDGEMENTS

My research has been supported out of my own pocket and by grants from the National Geographic Society, the John Simon

Guggenheim Memorial Foundation, a senior Fulbright Research Scholarship, the Australian-American Educational Foundation, the University Research Institute of the Graduate School at The University of Texas at Austin, the Denton A. Cooley Centennial Professorship in Zoology at The University of Texas at Austin, the U. S. National Science Foundation, and the U. S. National Aeronautics and Space Administration. I thank the staffs of the Department of Zoology at the University of Western Australia, the Western Australian Museum, the Department of Conservation and Land Management, and the Western Australian Department of Environment and Conservation. Lizards were collected under permits issued by the Department of Conservation and Land Management. Stephen Goodyear and Cyrus Harp helped with laboratory analyses of stomach contents. Scott Thompson and John Dell made useful comments and suggestions for improvements on the manuscript.

#### REFERENCES

- BAMFORD, M. J. 1998. Field observations on the semi-arboreal pygopodids *Aclys concinna* (Kluge 1974) and *Pletholax gracilis* (Cope 1864). *Herpetofauna - Sydney*, 28: 2-4.
- BAUER, A. M. 1986. Saltation in the Pygopodid Lizard, *Delma tincta*. *Journal of Herpetology*, 20: 462-463.
- BRADSHAW, S. D., GANS, C. and SAINT GIRONS, H. 1980. Behavioral Thermoregulation in a Pygopodid Lizard, *Lialis burtonis*. *Copeia*, 1980: 738-743.
- DOWNES, S., FOSTER, R., and MOLNAR, C. 1997. New insights into the distribution and habitat of the vulnerable bronzeback legless lizard *Ophidiocephalus taeniatus*. *Herpetofauna*, 27: 11-13.
- EHMANN, H. 1981. The natural history and conservation of the bronzeback (*Ophidiocephalus taeniatus* Lucas and Frost) (Lacertilia, Pygopodidae). In *Proceedings of the Melbourne Herpetological Symposium* (eds C.B. Brown and A.A. Martin.) pp. 7-13. (Zoological Board of Victoria, Melbourne.)
- EHMANN, H. 1992. The apparent severe decline of the bronzeback legless lizard (*Ophidiocephalus taeniatus*) at Abminga. *Herpetofauna*, 22: 31-33.
- EHMANN, H., and METCALFE, D. (1978). The rediscovery of *Ophidiocephalus taeniatus* Lucas and Frost (Pygopodidae, Lacertilia) the Bronzeback. *Herpetofauna*, 9: 8-10.
- HUEY, R. B. and SLATKIN, M. 1976. Costs and benefits of lizard thermoregulation. *Quarterly Review of Biology*, 51: 363-384.
- JENNINGS, W. B., PIANKA, E. R., and DONNELLAN, S. 2003. Systematics of the lizard family Pygopodidae with implications for the diversification of Australian temperate biotas. *Systematic Biology*, 52: 757-780.
- KLUGE, A. G. 1974. A taxonomic revision of the lizard family

- Pygopodidae. *Misc. Publ. Mus. Zool. Univ. Mich.* 147: 1-221.
- KLUGE, A. G. 1976. Phylogenetic relationships in the lizard family Pygopodidae: An evaluation of theory, methods and data. *Misc. Publ. Mus. Zool. Univ. Mich.* 152: 1-72.
- MURRAY, B. A., BRADSHAW, S. D. and EDWARD, D. H. 1991. Feeding Behavior and the Occurrence of Caudal Luring in Burton's Pygopodid *Lialis burtonis* (Sauria: Pygopodidae) *Copeia*, 1991: 509-516.
- NEILL, W. T. 1957. Notes on the Pygopodid Lizards, *Lialis burtonis* and *L. jicari*. *Copeia*, 1957: 230-232.
- PARKER, H. W. 1956. The lizard genus *Aprasia*; its taxonomy and temperature correlated variation. *Bull. Brit. Mus. Nat. Hist.* 3: 363-385.
- PATCHELL, F. C. and SHINE, R. 1986. Feeding Mechanisms in Pygopodid Lizards: How Can *Lialis* Swallow Such Large Prey? *Journal of Herpetology*, 20: 59-64.
- PATCHELL, F. C. and SHINE, R. 1986. Food Habits and Reproductive Biology of the Australian Legless Lizards (Pygopodidae). *Copeia*, 1986: 30-39.
- PATCHELL, F. C. and SHINE, R. 2009. Hinged teeth for hard-bodied prey: a case of convergent evolution between snakes and legless lizards. *Journal of Zoology*, 2082, 269-275.
- PIANKA, E. R. 1986. *Ecology and Natural History of Desert Lizards. Analyses of the Ecological Niche and Community Structure*. Princeton University Press, Princeton, New Jersey.
- SAVITZSKY, A. H. 1981. Hinged teeth in snakes: An adaptation for swallowing hard-bodied prey. *Science*, 212: 346-349
- SHINE, R. 1986. Evolutionary Advantages of Limblessness: Evidence from the Pygopodid Lizards. *Copeia*, 1986: 525-529.
- STEPHENSON, N. G. 1962. The comparative morphology of the head skeleton, girdles and hind limbs in the Pygopodidae. *Zoological Journal Linnean Society*, 44: 627-644.
- WEBB, J. K. and SHINE, R. 1994. Feeding Habits and Reproductive Biology of Australian Pygopodid Lizards of the Genus *Aprasia*. *Copeia*, 1994: 390-398.